

Geometric sequence

The n^{th} term of a
geometric sequence

$$u_n = u_1 r^{n-1}$$

The sum of n terms of a
finite geometric sequence

$$S_n = \frac{u_1(r^n - 1)}{r - 1} = \frac{u_1(1 - r^n)}{1 - r}, r \neq 1$$

The sum of an infinite
geometric sequence

$$S_\infty = \frac{u_1}{1 - r}, |r| < 1$$

Examples of geometric sequence

2, 10, 50, 250

1, 3, 9, 27, 81

5, -1, $\frac{1}{5}$, $-\frac{1}{25}$

u_n is the n^{th} term

r is the common ratio

$$r = \frac{u_2}{u_1} \text{ or } \frac{u_{n+1}}{u_n}$$

S_n is sum of n terms

Show geometric sequence:

$$\frac{u_2}{u_1} = \frac{u_3}{u_2}$$

Show Geometric sequence

1. Show that $12, -6, 3, -\frac{3}{2}, \dots$ is geometric and find the common ratio.

2. Show that $8, 4\sqrt{2}, 4, 2\sqrt{2}, \dots$ is geometric and find the common ratio.

List the terms

1. Consider the sequence defined by $u_n = 3(2)^{n-1}$

List the first four terms of the sequence.

2. Consider the sequence defined by $u_n = 4(-3)^{n-1}$

List the first four terms of the sequence.


Find the general term

$$u_n = u_1 r^{n-1}$$


1. A geometric sequence has $u_2 = -2$ and $u_7 = 64$. Find the expression of general term.


2. A geometric sequence has $u_3 = 8$ and $u_6 = -1$. Find the expression of general term.

Paper 1 exercise

1.  Three consecutive terms of a geometric sequence are $x - 3$,
6 and $x + 2$.

Find the possible values of x .


3.  The following diagram shows $[AB]$, with length 2 cm. The line is divided into an infinite number of line segments. The diagram shows the first three segments.



The length of the line segments are p cm, p^2 cm, p^3 cm, ... , where $0 < p < 1$.


Show that $p = \frac{2}{3}$.

(b) The following diagram shows $[CD]$, with length b cm, where $b > 1$. Squares with side lengths k cm, k^2 cm, k^3 cm, ... , where $0 < k < 1$, are drawn along $[CD]$. This process is carried on indefinitely. The diagram shows the first three squares.



The total sum of the areas of all the squares is $\frac{9}{16}$. Find the value of b .

Paper 2 exercise

1.  Consider a geometric sequence where the first term is 768 and the second term is 576. Find the least value of n such that the n^{th} term of the sequence is less than 7.

2.  (a) Consider an infinite geometric sequence with $u_1 = 40$ and

$$r = \frac{1}{2}.$$

(i) Find u_{14}


(ii) Find the sum of the infinite sequence.

Consider an arithmetic sequence with n terms, with first term (-36) and eighth term (-8) .

(b) (i) Find the common difference.

(ii) Show that $S_n = 2n^2 - 38n$.

(c) The sum of the infinite geometric sequence is equal to twice the sum of the arithmetic sequence. Find n .

3.  The diagrams below show the first four squares in a sequence of squares which are subdivided in half. The area of the shaded square is $\frac{1}{4}$.




Diagram 1




Diagram 2




Diagram 3




Diagram 4

- (a) (i) Find the area of square B and of square C.
 (ii) Show that the areas of squares A, B and C are in geometric progression.
 (iii) Write down the common ratio of the progression.
- (b) (i) Find the total area shaded in diagram 2.
 (ii) Find the total area shaded in the 8th diagram of this sequence.
 Give your answer correct to six significant figures.
- (c) The dividing and shading process illustrated is continued indefinitely. Find the total area shaded.

