


## Laws of probability

## $P(A \cup B) = P(A) + P(B) - P(A \cap B)$



1. If P(A) = 0.6, P(B) = 0.4 and  $P(A \cup B) = 0.7$ , find  $P(A \cap B)$ .

2. If P(A) = 0.5, P(B) = 0.4 and  $P(A \cup B) = 0.8$ , find  $P(A \cap B)$ .



### **Independent event**

$$P(A \cap B) = P(A) P(B)$$

The occurrence of each one of them does not affect the probability that the other occurs.

E.g. The probability of getting number 1 faces down in tossing a dice is independent to the probability of getting a head in throwing a fair coin.

1. If $P(A) = 0.5$ , $P(B) = 0.4$ and $P(A \cup B) = 0.7$ , are A and B	
independent events?	




3

### Mutually exclusive

$$P(A \cap B) = 0$$

$$P(A \cap B) = P(A) + P(B)$$



1. Given that P(A) = 0.5, P(B) = 0.3 and  $P(A \cup B) = k$ . Find k if (a) A and B are independent

(b) A and B are mutually exclusive



#### **Exercise**

## Paper 1

- 1. Events A and B are independent with  $P(A \cap B) = 0.2$  and  $P(A' \cap B) = 0.6$ .
- (a) Find (B).

#### IBDP Mathematics (SL) Laws of probability



- 2. Two events A and B are such that P(A) = 0.2 and  $P(A \cup B) = 0.5$ .
- (a) Given that A and B are mutually exclusive, find P(B).

(b) Given that A and B are independent, find P(B).



# Paper 2

1. Two events A and B are such that $P(A) = 0.2$ and $P(A \cup B) = 0.5$ .
(a) Given that $A$ and $B$ are mutually exclusive, find $P(B)$ .
(b) Given that $A$ and $B$ are independent, find $P(B)$ .

#### IBDP Mathematics (SL) Laws of probability



2. Et C and D be independent events, with P(C) = 2k and  $P(D) = 3k^2$ , where 0 < k < 0.5.

(a) Write down an expression for $P(C \cap D)$ in terms of $k$ .
(b) Given that $P(C \cap D) = 0.162$ , find $k$ .
(c) Find P(C' D).